

# Weighted (Co)Limits

David Jaz Myers

July 26, 2017

# Acknowledgements

- Thanks to the organizers of CT2017!
- Thanks to Emily, Alexander, and Brendan for running the Kan Extension Seminar!
- And thanks to all my fellow students in the seminar for their conversation and insight!

**Goal:** We want to develop a good theory of limits and colimits for enriched categories.

And give a few nice examples.

# Review of (Co)Limits

- We have a diagram shape (small category)  $\mathcal{D}$ , and an environment category  $\mathcal{E}$ .
- A *diagram* is a functor  $D : \mathcal{D} \rightarrow \mathcal{E}$ .
- A *cone* is a natural transformation  $1 \rightarrow \mathcal{E}(E, D(-))$  in  $\mathbf{Set}^{\mathcal{D}}$ .
- The *limit*  $\lim D$  of  $D$  is a universal cone:

$$\mathcal{E}(E, \lim D) \cong \mathbf{Set}^{\mathcal{D}}(1, \mathcal{E}(E, D(-))).$$

- For colimits, switch the direction of the cone around.

$$\mathcal{E}(\operatorname{colim} D, E) \cong \mathbf{Set}^{\mathcal{D}^{\text{op}}}(1, \mathcal{E}(D(-), E)).$$

# Thickening the Cones

- When sets are our base, we can analyze any set of morphisms in a category one at a time. That is, it suffices to look at points  $1 \rightarrow \mathcal{E}(E, D_i)$  in our cones.
- But in more general base categories, this is no longer the case. We need to be explicit about the shape of the legs of our cones.
- We will let the shape of the legs of our cones vary with the objects of the diagrams. These shapes are called *weights*.

## Definition

Let  $D : \mathcal{D} \rightarrow \mathcal{E}$  be a diagram in a category  $\mathcal{E}$  enriched in  $\mathcal{V}$ . Given a *functor of weights*  $W : \mathcal{D} \rightarrow \mathcal{V}$ , the **weighted limit**  $\lim_W D$ , if it exists, satisfies the following universal property:

$$\mathcal{E}(E, \lim_W D) \cong \mathcal{V}^{\mathcal{D}}(W(-), \mathcal{E}(E, D(-))).$$

# Four Examples

- ① Powers (over any base).
- ② Kernel Pairs (over sets).
- ③ Limits of Cauchy Sequences (over positive real numbers).
- ④ Homotopy Pushouts (over topological spaces).

- Let  $\mathcal{D} = 1$  be the walking object.
- A diagram  $D : \mathcal{D} \rightarrow \mathcal{E}$  is just an object of  $\mathcal{E}$ .
- A weight  $W : \mathcal{D} \rightarrow \mathcal{V}$  is just an object of the base.
- The weighted limit  $\lim_W D$  is given by

$$\mathcal{E}(E, \lim_W D) \cong \mathcal{V}(W, \mathcal{E}(E, D)),$$

showing that maps into  $\lim_W D$  are  $W$ -tuples of maps into  $D$ .  
Therefore,  $\lim_W D = D^W$ , the  $W$ -power of  $D$ .

# Kernel Pairs

- Let  $\mathcal{D} = \bullet \rightarrow \bullet$  be the walking arrow.
- A diagram  $D : \mathcal{D} \rightarrow \mathcal{E}$  is an arrow  $A \xrightarrow{f} B$  in  $\mathcal{E}$ .
- Take  $W : \mathcal{D} \rightarrow \mathbf{Set}$  to be  $2 \xrightarrow{!} 1$ .
- The weighted limit is then given by

$$\mathcal{E}(E, \lim_W D) \cong \mathbf{Set}^{\bullet \rightarrow \bullet}(2 \rightarrow 1, \mathcal{E}(E, A) \xrightarrow{f_*} \mathcal{E}(E, B)).$$

- Substituting in  $\lim_W D$  for  $E$  and pushing  $\mathbf{id}_{\lim_W D}$  through the isomorphism gives us

$$\begin{array}{ccc} 2 & \longrightarrow & \mathcal{E}(E, A) \\ \downarrow ! & & \downarrow f_* \\ 1 & \longrightarrow & \mathcal{E}(E, B) \end{array},$$

or, in  $\mathcal{E}$ ,

$$\lim_W D \rightrightarrows A \rightarrow B,$$

showing that  $\lim_W D$  is the kernel pair.



# Limits of Cauchy Sequences

- Let  $\mathcal{D} = \{0, 1, 2, \dots, \infty\}$  be the natural numbers, with  $\mathcal{D}(i, j) = \infty$ .
- A diagram  $D : \mathcal{D} \rightarrow \mathcal{E}$  is a sequence in  $\mathcal{E}$ .
- Let  $W : \mathcal{D} \rightarrow [0, \infty]$  be  $i \mapsto \frac{1}{2^i}$ , with  $\infty \mapsto 0$ .
- The weighted limit is then given by

$$\mathcal{E}(E, \lim_W D) = [0, \infty]^{\mathcal{D}}(W(-), \mathcal{E}(E, D(-))),$$

which means

$$0 = \mathcal{E}(\lim_W D, \lim_W D) = \sup_{i \in \mathcal{D}} \left( \mathcal{E}(E, D_i) - \frac{1}{2^i} \right),$$

so that  $D_i \rightarrow \lim_W D$  as a sequence.

# Homotopy Pushouts

- Let  $\mathcal{D} = \bullet \leftarrow \bullet \rightarrow \bullet$ , so that a diagram is a span in  $\mathcal{E}$ .
- Let  $W : \mathcal{D}^{\text{op}} \rightarrow \mathbf{Top}$  be the cospan  $* \xrightarrow{0} [0, 1] \xleftarrow{1} *$ .
- The weighted colimit is given by

$$\mathcal{E}(\text{colim}_W D, E) \cong \mathbf{Top}^{\mathcal{D}^{\text{op}}}(W(-), \mathcal{E}(D(-), E)),$$

which makes it the initial such cocone:

$$\begin{array}{ccc} C & \xrightarrow{i} & B \\ j \downarrow & \searrow & \downarrow \\ A & \longrightarrow & \text{colim}_W D \end{array}$$

Thanks for Listening!

Thanks for Listening!